

Math 308 Spring '18
Midterm 1
2018-04-20

Name: _____
Student ID Number: _____

Question	Points	Score
1	12	
2	12	
3	12	
4	12	
5	12	
Total:	60	

- There are 5 problems on this exam. Be sure you have all 5 problems on your exam.
- The final answer must be left in exact form. Box your final answer.
- You are allowed the TI-30XIIS calculator. It is possible to complete the exam without a calculator.
- You are allowed a single sheet of 2-sided handwritten self-written notes.
- You must show your work to receive full credit. A correct answer with no supporting work will receive a zero.
- Use the backsides if you need extra space. Make a note of this if you do.
- Do not cheat. This exam should represent your own work. If you are caught cheating, I will report you to the Community Standards and Student Conduct office.

Conventions:

- I will often denote the zero vector by 0 .
- When I define a variable, it is defined for that whole question. The A defined in Question 2 is the same for the entire page.
- I often use x to denote the vector (x_1, x_2, \dots, x_n) . It should be clear from context.
- Sometimes I write vectors as a row and sometimes as a column. The following are the same to me.

$$(1, 2, 3) \quad \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}.$$

- I write the evaluation of linear transforms in a few ways. The following are the same to me.

$$T(1, 2, 3) \quad T((1, 2, 3)) \quad T\left(\begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}\right)$$

2. Let

$$A = \begin{pmatrix} 1 & 1 & 0 & 3 \\ 0 & 1 & 1 & 1 \\ 1 & 2 & 1 & 4 \end{pmatrix} \quad \text{and} \quad b = \begin{pmatrix} 2 \\ -1 \\ 1 \end{pmatrix}$$

- (a) (6 points) Give the general solution to $Ax = b$. (Hint: The sum of the components in the reduced echelon form of A is 5.)
- (b) (3 points) Give 3 particular solutions to $Ax = b$ that do not lie on a line. To save time, you may consider naming some vectors. You may also leave your answer as a linear combination of vectors. (Here we are thinking of vectors as points in space.)
- (c) (3 points) Does there exist a vector $y \in \mathbb{R}^3$ such that $Ax = y$ has no solutions? Justify your answer. This is a yes-no question.

3. Let A and B be equivalent matrices given by

$$A = \begin{pmatrix} 1 & -1 & 3 & 6 \\ 2 & -2 & 0 & 6 \\ 3 & -3 & 8 & 17 \end{pmatrix} \sim \begin{pmatrix} 1 & -1 & 0 & 3 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 0 \end{pmatrix} = B.$$

Let $S = \{a_1, a_2, a_3, a_4\}$ be the set of columns of A . Let $T : \mathbb{R}^4 \rightarrow \mathbb{R}^3$ be a linear transformation defined by $T(x) = Ax$.

The following are quick questions that you should briefly justify with a single sentence or two.

(a) (2 points) Is T one-to-one?

(b) (2 points) Is T onto?

(c) (2 points) Is S linearly independent?

(d) (2 points) Is S spanning?

(e) (2 points) Let $b = 4a_1 + 20a_2 + 2018a_3$. What is the dimension of the solution space to $Ax = b$? (Hint: If you claim a solution exists or does not exist, you should explain why.)

(f) (2 points) Is there a choice of $b \in \mathbb{R}^3$ such that the solution space to $Ax = b$ contains finitely many points? (Same hint as above.)

4. Let

$$u_1 = \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}, u_2 = \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix}, u_3 = \begin{pmatrix} 3 \\ 2 \\ 1 \end{pmatrix}.$$

Let $T : \mathbb{R}^3 \rightarrow \mathbb{R}^2$ be the linear transformation such that

$$T(u_1) = \begin{pmatrix} 1 \\ 0 \end{pmatrix}, T(u_2) = \begin{pmatrix} 0 \\ 1 \end{pmatrix}, T(u_3) = \begin{pmatrix} 1 \\ 1 \end{pmatrix}.$$

(a) (4 points) Is T one-to-one? Is it onto? Briefly justify.

(b) (4 points) Write $e_3 = (0, 0, 1)$ as a linear combination of u_1, u_2, u_3 .

(c) (4 points) There is a matrix B such that $T(x) = Bx$. What is B ?

5. Let A and B be equivalent matrices given by

$$A = \begin{pmatrix} 1 & 3 & 5 & 4 & 13 \\ -1 & 0 & -2 & 2 & -1 \\ 1 & 8 & 10 & 0 & 19 \end{pmatrix} \sim \begin{pmatrix} 1 & 0 & 2 & 0 & 3 \\ 0 & 1 & 1 & 0 & 2 \\ 0 & 0 & 0 & 1 & 1 \end{pmatrix} = B$$

As usual, let a_1, a_2, a_3, a_4, a_5 denote the columns of A .

(a) (4 points) Give the homogeneous solution to $Ax = 0$.

(b) (4 points) Give the general solution to $2Ax = a_2 - a_4$. (Hint: $Ae_i = a_i$.)

(c) (4 points) The columns corresponding to free variables are a_3 and a_5 . Write them as a linear combination of a_1, a_2, a_4 .