

Math 308H - Winter 2018
Final
2018-03-15

Name: _____

Student ID Number: _____

Question	Points	Score
1	12	
2	12	
3	12	
4	12	
5	12	
6	12	
Total:	72	

- There are 6 problems on this exam. Be sure you have all 6 problems on your exam.
- The final answer must be left in exact form. Box your final answer.
- You are allowed the TI-30XIIS calculator. It is possible to complete the exam without a calculator.
- You are allowed a single sheet of 2-sided handwritten self-written notes.
- You must show your work to receive full credit. A correct answer with no supporting work will receive a zero.
- Use the backsides if you need extra space. Make a note of this if you do.
- Do not cheat. This exam should represent your own work. If you are caught cheating, I will report you to the Community Standards and Student Conduct office.

Conventions:

- I will often denote the zero vector by 0 .
- When I define a variable, it is defined for that whole question. The A defined in Question 2 is the same for each part.
- I treat row and column vectors as the same.
- For any linear transformation T , there exists a matrix A such that $T(x) = Ax$. I defined the determinant, rank, and nullity of T using A . This means,

$$\det(T) = \det(A), \quad \text{rank}(T) = \text{rank}(A), \quad \text{nullity}(T) = \text{nullity}(A).$$

3. Let A and B be equivalent matrices defined by

$$A = \begin{bmatrix} 1 & 2 & 0 & -1 \\ 0 & 1 & 0 & 3 \\ 0 & 1 & 1 & 0 \\ 2 & 5 & 0 & 1 \end{bmatrix} \sim \begin{bmatrix} 1 & 0 & 0 & -7 \\ 0 & 1 & 0 & 3 \\ 0 & 0 & 1 & -3 \\ 0 & 0 & 0 & 0 \end{bmatrix} = B$$

Let a_1, a_2, a_3, a_4 denote the columns of A .

(a) (3 points) Do not write express a basis as a matrix.

1. Give a basis for $\text{col}(2A^t)$.

2. Give a basis for $\text{null}(A)$.

3. Give a basis for $\text{row}(A)$.

(b) (3 points) These should be quick questions.

1. What is $\text{rank}(A)$?

2. What is $\text{nullity}(A^t D^{-1})$, where D is the 4×4 diagonal matrix consisting of 1, 2, 3, 4 along the diagonal.

3. What is $\det(2A)$?

(c) (3 points) Give a nontrivial linear combination of the columns of A that sum to zero. You may use a_1, a_2, a_3, a_4 to denote the columns of A .

(d) (3 points) Let C be the 4×3 matrix given by $C = [a_1 \ a_2 \ a_3]$. So C is the submatrix of A consisting of the first 3 columns. Give the general solution for $Cx = a_1 + a_4$.

4. Let $T : \mathbb{R}^4 \rightarrow \mathbb{R}^3$ be the linear transformation defined by

$$T(w, x, y, z) = (w + y + z, x + y + z, x + y + z).$$

(a) (3 points) There is a matrix A such that $T(x) = Ax$. What is A ?

(b) (3 points) Let $v = (0, 3, 0, 8)$. Give the general solution to $Ax = 2Av + (2, 1, 1)$.

(c) (3 points) Does there exist a rank 2 linear transformation S such that $T \circ S$ is the zero transformation? If so, give an example. If not, why not?

(d) (3 points) Does there exist a rank 3 linear transformation S such that $T \circ S$ is the zero transformation? If so, give an example. If not, why not?

5. Let

$$A = \begin{bmatrix} 0 & -1 & \frac{37}{3} & -\frac{253}{15} \\ 0 & 2 & 0 & -\frac{1}{5} \\ 0 & 0 & 2 & \frac{7}{5} \\ 0 & 0 & 0 & 3 \end{bmatrix}$$

be a matrix which decomposes as $A = UDU^{-1}$, where

$$U = \begin{bmatrix} 1 & -1 & 18 & 1 \\ 0 & 2 & 1 & -1 \\ 0 & 0 & 3 & 7 \\ 0 & 0 & 0 & 5 \end{bmatrix}, \quad D = \begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & 2 & 0 & 0 \\ 0 & 0 & 2 & 0 \\ 0 & 0 & 0 & 3 \end{bmatrix}.$$

Let u_1, u_2, u_3, u_4 be the columns of U and $\mathcal{B} = \{u_1, u_2, u_3, u_4\}$.

(a) (6 points) Fill out this table.

Eigenvalue λ	Alg. Multiplicity of λ	Geo. Multiplicity of λ	Basis for E_λ

(b) (3 points) Let $x = u_1 + u_2 + u_3 + u_4$. Express $A^{18}x$ as a linear combination of u_1, u_2, u_3, u_4 . You are allowed to have exponents of numbers in your answer. (Hint: x has been expressed as the sum of eigenvectors.)

(c) (3 points) What are the eigenvalues for $A^2 - 2A$?

6. Let $T(x) = Ax$, where A is as defined in Question 5. Let u_1, u_2, u_3, u_4 also be as defined in Question 5.
- (a) (4 points) Give two vectors v, w such that the triangle with vertices $\{T(0), T(v), T(w)\}$ has 6 times the area as the triangle with vertices $\{0, v, w\}$. Be sure to justify your answer. (Hint: It is unnecessary to compute the area of these triangles.)

- (b) (4 points) Find a basis for each of the following subspaces. If a subspace is trivial, then write \emptyset for its basis.

- $\text{null}(A - 2I)$

- $\text{null}(A^2 - 3I)$.

- (c) (4 points) Let $B = \{u_1, u_2, u_3, u_4\}$ be a basis.

- What is the general solution to $Ax = u_2 + 2u_3$?

- Let y be a particular solution to the above linear system. What is $[y]_B$?