


Lecture #16


16.6 (cont.)

Recall from 15.5 (text) : Surface integrals



we will think of this as a "uv"-plane when working with $\vec{r}(u,v)$, i.e. parametric surfaces.

Computing a small piece of Surface area:



We estimate small pieces by the area of a parallelogram!

To do this, we took ^{the magnitude of the} cross product of $|\vec{a} \times \vec{b}|$, since this gives us the area of the parallelogram spanned by \vec{a} and \vec{b} .


From last lecture, recall

\vec{r}_u and \vec{r}_v ("partials in the direction of u and v on a parametric surface")

We can use these in the place of \vec{a} and \vec{b} !

If you look back to the lecture notes on Surface area,

You will see that this is secretly what we were using all along, but instead of u, v , we used x and y , and constrained ourselves to usual surfaces (not parametric ones!) So,



Computing:

$$\text{Area}(\Delta S_{ij}) \approx \text{Area}(\Delta P_{ij}) = |\Delta u \vec{r}_u \times \Delta v \vec{r}_v| = |\vec{r}_u \times \vec{r}_v| \Delta u \Delta v.$$

Summing over each piece in the surface:

$$\sum_{i=1}^m \sum_{j=1}^n |\vec{r}_{u,ij}^* \times \vec{r}_{v,ij}^*| \Delta u \Delta v$$

and taking a limit, we get:

DEF If a smooth parametric surface S is given by the equation

$$\vec{r}(u, v) = x(u, v) \hat{i} + y(u, v) \hat{j} + z(u, v) \hat{k}, \quad (u, v) \in D$$

and S is covered just once as (u, v) ranges throughout the parameter domain D , then the surface area of S is

$$A(S) = \iint_D |\vec{r}_u \times \vec{r}_v| dA$$

where $\vec{r}_u = \frac{\partial x}{\partial u} \hat{i} + \frac{\partial y}{\partial u} \hat{j} + \frac{\partial z}{\partial u} \hat{k}$ and

$$\vec{r}_v = \frac{\partial x}{\partial v} \hat{i} + \frac{\partial y}{\partial v} \hat{j} + \frac{\partial z}{\partial v} \hat{k}$$

EXAMPLE 9

Find the surface area of a sphere of radius a .

rem: Parametric equation of a sphere:

$$x = a \sin \phi \cos \theta$$

$$y = a \sin \phi \sin \theta$$

$$z = a \cos \phi,$$

→ [For $0 \leq \phi \leq \pi$ and $0 \leq \theta \leq 2\pi$.]

This is the parameter domain D;

$$D = \{(\phi, \theta) : 0 \leq \phi \leq \pi, 0 \leq \theta \leq 2\pi\}$$

We start by computing $\vec{\Gamma}_\phi \times \vec{\Gamma}_\theta$

$$\vec{\Gamma}_\phi \times \vec{\Gamma}_\theta = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ \frac{\partial x}{\partial \phi} & \frac{\partial y}{\partial \phi} & \frac{\partial z}{\partial \phi} \\ \frac{\partial x}{\partial \theta} & \frac{\partial y}{\partial \theta} & \frac{\partial z}{\partial \theta} \end{vmatrix} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ a \cos \phi \cos \theta & a \cos \phi \sin \theta & -a \sin \phi \\ -a \sin \phi \sin \theta & a \sin \phi \cos \theta & 0 \end{vmatrix}$$

$$= a^2 \sin^2 \phi \cos \theta \hat{i} + a^2 \sin^2 \phi \sin \theta \hat{j} + (a^2 \cos \phi \cos^2 \theta \sin \phi + a^2 \cos \phi \sin^2 \theta \sin \phi) \hat{k}$$

$$= a^2 \sin^2 \phi \cos \theta \hat{i} + a^2 \sin^2 \phi \sin \theta \hat{j} + (a^2 \cos \phi \sin \phi) \hat{k}$$

Then: $|\vec{\Gamma}_\phi \times \vec{\Gamma}_\theta| = \sqrt{a^4 \sin^4 \phi \cos^2 \theta + a^4 \sin^4 \phi \sin^2 \theta + a^4 \cos^2 \phi \sin^2 \phi}$

$$= \sqrt{a^4 \sin^4 \phi + a^4 \cos^2 \phi \sin^2 \phi}$$

$$= a^2 \sqrt{\sin^4 \phi + \cos^2 \phi \sin^2 \phi}$$

$$= a^2 \sqrt{\sin^4 \phi + (1 - \sin^2 \phi) \sin^2 \phi}$$

$$= a^2 \sqrt{\sin^4 \phi + \sin^2 \phi - \sin^4 \phi}$$

$$= a^2 \sqrt{\sin^2 \phi}$$

$$= \boxed{a^2 \sin \phi}$$

* What does this remind you of? (Jacobians of sorts...?)

Then
$$\iint_D |\vec{r}_\phi \times \vec{r}_\theta| dA = \int_0^{2\pi} \int_0^\pi a^2 \sin\phi d\phi d\theta$$

$$= a^2 \int_0^{2\pi} [-\cos\phi]_0^\pi d\theta$$

$$= a^2 \int_0^{2\pi} 2 d\theta$$

$$= a^2 [2\theta]_0^{2\pi}$$

$$= a^2 \cdot 4\pi \quad \Rightarrow \quad \boxed{= 4\pi a^2}$$

EXERCISE Let your parametrized surface be described by

$$\begin{cases} x=x \\ y=y \\ z=f(x,y) \end{cases}$$

Show

$$\boxed{A(s) = \iint_D \sqrt{1 + \left(\frac{\partial z}{\partial x}\right)^2 + \left(\frac{\partial z}{\partial y}\right)^2} dA}$$

REMARK

This is telling you that the graph of a function is a special case of a parametric surface, and our new equation for surface area agrees with our old one!

↙ If you want to brush up on using this formula

EXERCISE

Find the area of the part of the paraboloid $z = x^2 + y^2$ that lies under the plane $z = 9$.

16.7 Surface Integrals

REMARK Compare what follows to our treatment of line integrals in 16.2. You'll find it analogous in many respects!

DEF We define a surface integral of f over the surface S as

$$\iint_S f(x,y,z) dS = \lim_{m,n \rightarrow \infty} \sum_{j=1}^m \sum_{i=1}^n f(P_{ij}^*) \Delta S_{ij} = \lim_{m,n \rightarrow \infty} \sum_{j=1}^m \sum_{i=1}^n f(P_{ij}^*) |\vec{r}_u \times \vec{r}_v| \Delta u \Delta v$$

← just add the function in!

Given a parametric equation $\vec{r}(u,v)$ describing the surface S , we can compute the surface integral as follows:

$$\boxed{\iint_S f(x,y,z) dS = \iint_D f(\vec{r}(u,v)) |\vec{r}_u \times \vec{r}_v| dA}$$

↑
parameter domain

REMARK If $f(x,y,z) = 1$, then $\iint_S f(x,y,z) dS = \iint_S dS = \iint_D |\vec{r}_u \times \vec{r}_v| dA = A(S)$
(surface area of S)

EXAMPLE 1 Compute $\iint_S x^2 dS$, where S is the unit sphere $x^2 + y^2 + z^2 = 1$.

STEP 1 Parametrize! $\vec{r}(\phi, \theta) = \langle \overset{x}{\sin \phi \cos \theta}, \overset{y}{\sin \phi \sin \theta}, \overset{z}{\cos \phi} \rangle$

$$D = \{(\phi, \theta) : 0 \leq \phi \leq \pi, 0 \leq \theta \leq 2\pi\}$$

STEP 2 compute $|\vec{r}_\phi \times \vec{r}_\theta|$. (We just did this! For radius a , we got $a^2 \sin \phi$. Here, we get...)

$$|\vec{r}_\phi \times \vec{r}_\theta| = \sin\phi$$

STEP 3. Compute

$$\iint_S x^2 dS = \iint_D \overset{x(\phi, \theta)}{\downarrow} (\sin\phi \cos\theta)^2 \cdot |\vec{r}_\phi \times \vec{r}_\theta| dA$$

$$= \int_0^{2\pi} \int_0^\pi \sin^3\phi \cos^2\theta d\phi d\theta$$

$$= \int_0^{2\pi} \cos^2\theta \left(\int_0^\pi (\sin\phi - \underbrace{\cos^2\phi \sin\phi}_{u\text{-sub}}) d\phi \right) d\theta$$

$\sin^2\phi - \sin\phi = (1 - \cos^2\phi) \sin\phi$

$$= \int_0^{2\pi} \cos^2\theta \left[-\cos\phi + \frac{\cos^3\phi}{3} \right]_0^\pi d\theta$$

$$= \int_0^{2\pi} \cos^2\theta \cdot \left(\left(1 - \frac{1}{3}\right) - \left(-1 + \frac{1}{3}\right) \right) d\theta$$

$$= \int_0^{2\pi} \cos^2\theta \cdot \left(\frac{4}{3}\right) d\theta$$

$$= \frac{4}{3} \int_0^{2\pi} \frac{1}{2} (1 + \cos 2\theta) d\theta$$

double angle identity!

$$= \frac{4}{3} \cdot \frac{1}{2} \left[\theta + \frac{1}{2} \sin 2\theta \right]_0^{2\pi}$$

$$= \frac{4}{3} \cdot \frac{1}{2} \left[(2\pi + 0) - (0 + 0) \right]$$

$$= \frac{4}{3} \cdot \frac{1}{2} \cdot 2\pi$$

$$= \frac{4}{3} \pi$$

REMARK We can use surface integrals to find the "mass" of parametric surfaces with density $\rho(x,y,z)$:

$$m = \iint_S \rho(x,y,z) dS$$

and the "center of mass" $(\bar{x}, \bar{y}, \bar{z})$:

$$\bar{x} = \frac{1}{m} \iint_S x \rho(x,y,z) dS, \quad \bar{y} = \frac{1}{m} \iint_S y \rho(x,y,z) dS, \quad \bar{z} = \frac{1}{m} \iint_S z \rho(x,y,z) dS$$

Now, let's switch to a special case of parametric surfaces, graphs of functions (i.e. just a surface $z = g(x,y)$).

Rem we can always think of a graph of a function as

$$\begin{cases} x = x \\ y = y \\ z = g(x,y) \end{cases},$$

a set of parametric equations.

" $\vec{r}(x,y)$ "

For this special case,

this is " $f(\vec{r}(x,y))$ "

$$\iint_S f(x,y,z) dS = \iint_D f(x, y, g(x,y)) \cdot |\vec{r}_x \times \vec{r}_y| dA$$

$$= \iint_D f(x, y, g(x,y)) \cdot \sqrt{1 + \left(\frac{\partial z}{\partial x}\right)^2 + \left(\frac{\partial z}{\partial y}\right)^2} dA$$

(*)

(*)

EXAMPLE 2 Evaluate $\iint_S y \, dS$, where S is the surface
 $z = x + y^2$, $0 \leq x \leq 1$, $0 \leq y \leq 2$.

STEP 1 : $\frac{\partial z}{\partial x} = 1$
 $\frac{\partial z}{\partial y} = 2y$

STEP 2 : Use our new formula!

$$\iint_S y \, dS = \iint_D y \cdot \sqrt{1 + \left(\frac{\partial z}{\partial x}\right)^2 + \left(\frac{\partial z}{\partial y}\right)^2} \, dA$$

$$= \int_0^1 \int_0^2 y \sqrt{1 + 1^2 + 4y^2} \, dy \, dx$$

$$= \int_0^1 \int_0^2 y \sqrt{2 + 4y^2} \, dy \, dx$$

$$= \int_0^1 \left. \frac{1}{8} \right|_{y=0}^{y=2} \sqrt{u} \, du \, dx$$

$$u = 2 + 4y^2, \quad du = 8y \, dy$$

$$= \int_0^1 \frac{1}{8} \cdot \left[\frac{2}{3} (2 + 4y^2)^{3/2} \right]_0^2 \, dx$$

$$= \int_0^1 \frac{1}{8} \cdot \left[\frac{2}{3} \left((18\sqrt{18}) - 2\sqrt{2} \right) \right] \, dx$$


$$= \frac{1}{8} \cdot \frac{2}{3} \cdot (54\sqrt{2} - 2\sqrt{2})$$

$$= \frac{1}{8} \cdot \frac{2}{3} \cdot 52\sqrt{2} = \frac{13\sqrt{2}}{3}$$

$52 = 13 \cdot 4$

REMARK

If a surface S is composed of a series of smaller surfaces, such as:



$$S = S_1 + S_2 + S_3$$

Then

$$\iint_S f(x, y, z) dS = \iint_{S_1} f(x, y, z) dS + \dots + \iint_{S_n} f(x, y, z) dS$$


EXAMPLE 3

Evaluate $\iint_S z dS$, where S is the surface whose sides

S_1 are given by the cylinder $x^2 + y^2 = 1$, whose bottom

is the disk $x^2 + y^2 \leq 1$ in the plane $z = 0$, and whose top S_3

is the part of the plane $z = 1 + x$ that lies above S_2 .



To integrate $\iint_S z \, dS$, we integrate each side separately

$$\iint_S z \, dS = \iint_{S_1} z \, dS + \iint_{S_2} z \, dS + \iint_{S_3} z \, dS$$

Starting with S_1 :

Step 1 parametrize S_1 , $x^2 + y^2 = 1$

$$\begin{cases} x = \cos \theta \\ y = \sin \theta \\ z = z \end{cases} \begin{cases} \rightarrow 0 \leq \theta \leq 2\pi \\ \rightarrow z \text{ goes from } 0 \text{ up to } 1+x \\ 0 \leq z \leq 1+x = \underline{1 + \cos \theta} \end{cases}$$

So:

$$\begin{cases} \vec{r}(\theta, z) = \cos \theta \hat{i} + \sin \theta \hat{j} + z \hat{k} \\ D = \{ (\theta, z) : 0 \leq \theta \leq 2\pi, 0 \leq z \leq 1 + \cos \theta \} \end{cases}$$

Step 2 compute $|\vec{r}_\theta \times \vec{r}_z|$

$$\vec{r}_\theta \times \vec{r}_z = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ -\sin \theta & \cos \theta & 0 \\ 0 & 0 & 1 \end{vmatrix} = \cos \theta \hat{i} + \sin \theta \hat{j} + 0 \hat{k}$$

$$|\vec{r}_\theta \times \vec{r}_z| = 1$$

STEP 3 compute

$$\begin{aligned} \iint_{S_1} z \, dS &= \int_0^{2\pi} \int_0^{1+\cos \theta} z |\vec{r}_\theta \times \vec{r}_z| \, dz \, d\theta = \int_0^{2\pi} \int_0^{1+\cos \theta} z \, dz \, d\theta \\ &= \int_0^{2\pi} \frac{(1+\cos \theta)^2}{2} \, d\theta \end{aligned}$$

$$\begin{aligned}
&= \frac{1}{2} \int_0^{2\pi} (1 + 2\cos\theta + \underbrace{\cos^2\theta}_{\downarrow \text{double angle!}}) d\theta \\
&= \frac{1}{2} \int_0^{2\pi} (1 + 2\cos\theta + \underbrace{\frac{1}{2}(1 + \cos 2\theta)}) d\theta \\
&= \frac{1}{2} \left[\theta + 2\sin\theta + \frac{1}{2}\theta + \frac{1}{4}\sin(2\theta) \right]_0^{2\pi}
\end{aligned}$$

$$= \boxed{\frac{3\pi}{2}}$$

Next, we compute S_2 . Notice that for S_2 , $\underline{z=0}$, so

$$\iint_{S_2} z dS = \iint_{S_2} 0 dS = 0$$

Next, S_3 . Notice S_3 is the graph of the function $f(x,y) = 1+x$,

above $x^2+y^2 \leq 1$. Thus, we know that $|\vec{r}_x \times \vec{r}_y| = \sqrt{1 + \left(\frac{\partial z}{\partial x}\right)^2 + \left(\frac{\partial z}{\partial y}\right)^2}$.

$$\iint_{S_3} z dS = \iint_D \overset{z=1+x}{(1+x)} \cdot \sqrt{1 + 1^2 + 0^2} dA, \quad D = \{(x,y) : x^2+y^2 \leq 1\}$$

↙ use polar!

$$= \int_0^{2\pi} \int_0^1 (1+r\cos\theta) \sqrt{2} \cdot r dr d\theta$$

$$= \sqrt{2} \int_0^{2\pi} \int_0^1 (r + r^2 \cos\theta) dr d\theta$$

$$= \sqrt{2} \int_0^{2\pi} \left[\frac{r^2}{2} + \frac{r^3}{3} \cos\theta \right]_0^1 d\theta$$

$$= \sqrt{2} \int_0^{2\pi} \left(\frac{1}{2} + \frac{1}{3} \cos\theta \right) d\theta$$

$$= \sqrt{2} \cdot \left[\frac{1}{2} \theta + \frac{1}{3} \sin \theta \right]_0^{2\pi}$$

$$= \sqrt{2} \cdot [(\pi + 0) - (0 + 0)]$$

$$= \boxed{\sqrt{2} \pi}$$

Thus, we can conclude that

$$\iint_S z \, dS = \iint_{S_1} z \, dS + \iint_{S_2} z \, dS + \iint_{S_3} z \, dS = \frac{3\pi}{2} + 0 + \sqrt{2} \pi$$

$$= \boxed{\frac{3\pi}{2} + \sqrt{2} \pi}$$

Exercises (16.6) #19, 23, 25, 33, 35, 40, 47, 49

and #61-63, and if feeling bold: 64.

Exercises (16.7) #8, 9, 17